



**Entresto**<sup>®</sup>  
(sacubitril/valsartan) tablets  
24/26mg • 49/51mg • 97/103mg



## HEADING HOME WITH ENTRESTO<sup>®</sup>

Follow the 4 steps below

### 1 Start

Start taking your ENTRESTO prescription as soon as advised by your doctor. See inside to learn how to save.

### 2 Follow up

Follow up within 7 to 14 days to lower the risk of hospitalization. Schedule an appointment and stay on top of it!

### 3 Prepare

Prepare for your doctor visit. Fill out the enclosed Doctor Discussion Guide and note any symptoms and questions you may have.

### 4 Prioritize

Prioritize Heart Failure. It's an ongoing condition that can get worse over time and needs your attention.

#### What is ENTRESTO?

ENTRESTO is a prescription medicine used to treat adults with long-lasting (chronic) heart failure to help reduce the risk of death and hospitalization. ENTRESTO works better when the heart cannot pump a normal amount of blood to the body.

#### IMPORTANT SAFETY INFORMATION

##### What is the most important information I should know about ENTRESTO?

ENTRESTO can harm or cause death to your unborn baby. Talk to your doctor about other ways to treat heart failure if you plan to become pregnant. Tell your doctor right away if you become pregnant during treatment with ENTRESTO.

See additional Important Safety Information throughout and on page 5. Please [click here](#) for full Prescribing Information.

# HOW TO TAKE ENTRESTO®

**ENTRESTO is a twice-daily medicine.** You should take your dose of ENTRESTO exactly as prescribed by your doctor. ENTRESTO is usually given with other Heart Failure medicines in place of an ACE inhibitor or other ARB. **Take ENTRESTO every day as prescribed. If you stop taking ENTRESTO, the medicine will stop working for you.**

**ENTRESTO is a tablet that comes in the following doses:**



**24/26 mg**



**49/51 mg**



**97/103 mg**

**If you have been prescribed ENTRESTO and are currently taking an ACE inhibitor\*:**



**Stop taking your  
ACE  
inhibitor\***



**Wait for  
36  
hours**



**Start taking  
ENTRESTO,  
as prescribed**

\*Such as enalapril or lisinopril.

## IMPORTANT SAFETY INFORMATION (cont)

**Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your doctor if you take potassium supplements or a salt substitute; nonsteroidal anti-inflammatory drugs (NSAIDs); lithium; or other medicines for high blood pressure or heart problems such as an ACE inhibitor, ARB, or aliskiren.

### Do not take ENTRESTO if you:

- are allergic to any of the ingredients in ENTRESTO
- have had an allergic reaction including swelling of your face, lips, tongue, throat (angioedema), or trouble breathing while taking a type of medicine called an angiotensin-converting enzyme (ACE) inhibitor or angiotensin II receptor blocker (ARB)

**See additional Important Safety Information throughout and on page 5. Please [click here](#) for full Prescribing Information.**



**Entresto®**  
(sacubitril/valsartan) tablets  
24/26 mg • 49/51 mg • 97/103 mg

# HELPFUL TOOLS FOR GETTING STARTED ON ENTRESTO®

**Enspire**  
PROGRAM  
from Entresto®  
SUPPORT DESIGNED WITH YOU IN MIND



## Sign up for free lifestyle and treatment support

The ENSPIRE Program from ENTRESTO® provides tips and tools to support you as you manage your Heart Failure, including resources that can help you keep track of your medications, stay active, and eat healthier.

## Ways to enroll in the ENSPIRE Program from ENTRESTO®



Call **1-888-368-7378**

(Monday–Friday, excluding holidays,  
8:00 AM to 8:00 PM ET)



**Click here** to enroll.

### Additional Resources

The resources below will help you understand what to expect with ENTRESTO and prepare for your next appointment.

- Download the **Doctor Discussion Guide** [here](#)
- Download the brochure “**Getting Started on ENTRESTO**” [here](#)

### IMPORTANT SAFETY INFORMATION (cont)

#### Do not take ENTRESTO if you (cont):

- take an ACE inhibitor medicine. Do not take ENTRESTO for at least 36 hours before or after you take an ACE inhibitor medicine. Talk with your doctor or pharmacist before taking ENTRESTO if you are not sure if you take an ACE inhibitor medicine
- have diabetes and take a medicine that contains aliskiren


See additional Important Safety Information throughout and on page 5. Please [click here](#) for full Prescribing Information.



**Entresto®**  
(sacubitril/valsartan) tablets  
24/26 mg • 49/51 mg • 97/103 mg

## 30-Day Free Trial Offer

For all patients (regardless of insurance) when initiating treatment. This voucher is good for a 30-day (maximum 60 tablets, one-time use) free trial of ENTRESTO® at no cost to you. One per patient. Limitations apply.



**Just bring these pre-activated codes to your pharmacy with a valid prescription to redeem!**

## \$10 Co-Pay Offer

- For eligible commercially/privately insured patients. Pay as little as \$10 for up to a 90-day fill
- To sign up call **1-888-368-7378** or **click here**

Once you sign up, your co-pay registration will automatically be renewed at the beginning of each calendar year (annual limit of \$4100). Not valid under Medicare, Medicaid, or any other federal or state program. Limitations apply. See Terms and Conditions [here](#).



**See patient stories, recipes, and helpful tips for living with Heart Failure on our YouTube channel and Facebook page**

 [Facebook.com/entresto](https://www.facebook.com/entresto)

 [@entresto\\_sacubitril\\_valsartan](https://www.instagram.com/entresto_sacubitril_valsartan)

 [YouTube.com/LivingWithHF](https://www.youtube.com/LivingWithHF)

**Visit our website to learn more about these offers and to enroll.**

The Novartis Patient Assistance Foundation, also known as NPAF, provides Novartis medicines at no cost to eligible US patients who have **limited or no prescription drug coverage** and cannot afford the cost of their medication. If eligible, Novartis medications may be available for free.

Call NPAF at **1-800-277-2254** or visit [www.PAP.Novartis.com](http://www.PAP.Novartis.com) for more information or to download an application.

### IMPORTANT SAFETY INFORMATION (cont)

**Before taking ENTRESTO tell your doctor about all of your medical conditions, including if you:**

- have a history of hereditary angioedema
- have kidney or liver problems
- have diabetes
- are pregnant or plan to become pregnant; are breastfeeding or plan to breastfeed. You should not breastfeed during treatment with ENTRESTO

**See additional Important Safety Information throughout and on page 5. Please [click here](#) for full Prescribing Information.**

 **Entresto®**  
(sacubitril/valsartan) tablets  
24/26mg • 49/51mg • 97/103mg

## What is ENTRESTO®?

ENTRESTO is a prescription medicine used to treat adults with long-lasting (chronic) heart failure to help reduce the risk of death and hospitalization. ENTRESTO works better when the heart cannot pump a normal amount of blood to the body.

## IMPORTANT SAFETY INFORMATION

### What is the most important information I should know about ENTRESTO?

ENTRESTO can harm or cause death to your unborn baby. Talk to your doctor about other ways to treat heart failure if you plan to become pregnant. Tell your doctor right away if you become pregnant during treatment with ENTRESTO.

### Do not take ENTRESTO if you:

- are allergic to any of the ingredients in ENTRESTO
- have had an allergic reaction including swelling of your face, lips, tongue, throat (angioedema), or trouble breathing while taking a type of medicine called an angiotensin-converting enzyme (ACE) inhibitor or angiotensin II receptor blocker (ARB)
- take an ACE inhibitor medicine. Do not take ENTRESTO for at least 36 hours before or after you take an ACE inhibitor medicine. Talk with your doctor or pharmacist before taking ENTRESTO if you are not sure if you take an ACE inhibitor medicine
- have diabetes and take a medicine that contains aliskiren

### Before taking ENTRESTO tell your doctor about all of your medical conditions, including if you:

- have a history of hereditary angioedema
- have kidney or liver problems
- have diabetes
- are pregnant or plan to become pregnant; are breastfeeding or plan to breastfeed. You should not breastfeed during treatment with ENTRESTO

**Tell your doctor about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your doctor if you take potassium supplements or a salt substitute; nonsteroidal anti-inflammatory drugs (NSAIDs); lithium; or other medicines for high blood pressure or heart problems such as an ACE inhibitor, ARB, or aliskiren.

### What are the possible side effects of ENTRESTO? ENTRESTO may cause serious side effects including:

- **swelling of your face, lips, tongue, and throat (angioedema) that may cause trouble breathing and death.** Get emergency medical help right away if you have symptoms of angioedema or trouble breathing. Do not take ENTRESTO again if you have had angioedema while taking ENTRESTO
- people who are Black or who have had angioedema and take ENTRESTO may have a higher risk of having angioedema
- **low blood pressure (hypotension),** which is common, and your risk of low blood pressure is greater if you also take water pills (diuretics). Call your doctor if you become dizzy or lightheaded, or you develop extreme tiredness (fatigue)
- **kidney problems,** which are common, and can be serious and can lead to kidney failure
- **increased amount of potassium in your blood (hyperkalemia),** which is common

The most common side effects also include cough and dizziness.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.



**Entresto®**  
(sacubitril/valsartan) tablets  
24/26mg • 49/51mg • 97/103mg

Please [click here](#) for full Prescribing Information, including Boxed WARNING.



Novartis Pharmaceuticals Corporation  
East Hanover, New Jersey 07936-1080

© 2024 Novartis

6/24

441424